

Jamia Hamdard
School of Management and Business Studies
Department of Management
Odd Semester Examination - January 2023
BBA/BBAMBA-Integrated- V Semester
Subject Code- BBA-F-17/BBA(I)-F-17
Subject Name- Quantitative Techniques for Managers

Roll No...

Duration: 3 Hour

Max. Marks -75

Section A - Short Answer

(5X6=30)

Note: Attempt any Six Questions; All questions carry equal marks

Q1: Differentiate between the term 'Sample' & 'Population'.

Q2: Define standard error of estimate. How is r^2 a better measure than the standard error of estimate.

Q3: A manufacturing firm produces pipes in two plants I and II with daily production of 1500 and 2000 pipes respectively. The fraction of defective pipes produced by two plants 1 and 2 are 0.006 and 0.008 respectively. If pipe selected at random from that day's production is found to be defective, what is the chance that it has come from plant I and plant II.

Q4: An aptitude test for selecting officers in a bank was conducted on 1000 candidates, the average score is 42 and the standard deviation of scores is 24. Assuming normal distribution for the scores, find:

a) The number of candidates whose score exceeds 58.

b) The number of candidates whose score lies between 30 and 66.

Q5: Define decision tree. Explain the different steps through which the decision maker proceeds in a decision analysis.

Q6: Distinguish between Risk and Uncertainty.

Q7: An unbiased coin is tossed twice, what is the probability of getting exactly 'two heads' & 'no heads'?

Q8: Explain the different applications of Chi-square test.

Q9: Define sampling frame.

Section B-Long Answer

(15X3=45)

Note: Attempt any Three Questions; All questions carry equal marks

Q1: What is sampling? Discuss different types of sampling techniques with example.

Q2: The following table gives monthly sales (in thousand rupees) of a certain firm in three states by its four salesmen:

States	Salesman			
	I	II	III	IV
A	6	5	3	8
B	8	9	6	5
C	10	7	8	7

Set up the analysis of variance table and test whether there is any significant difference

(i) between sales by the firm salesman, and

(ii) sales in the three states

Q3: (a) Two working designs are under consideration for adoption in a plant. A time and motion study shows that 12 workers using design 'A' have a mean assembly time of 300 seconds with a standard deviation of 12 seconds and that 15 workers using design 'B' have a mean assembly time of 335 seconds with a standard deviation of 15 seconds. Is the difference in the mean assembly time between the two working designs significant at one per cent level of significance?

(b) If 54 out of a random sample of 150 boys smoke, while 31 out of random sample of 100 girls smoke, can we conclude at 0.05 level of significance that the proportion of male smokers is higher than that of female smokers?

Q4: Explain the various steps involved in hypothesis testing. When do we use Z test & T test?

Q5: From the data given below find:

(a) the two regression equations

(b) the coefficient of correlation between marks in Economics and Statistics

Marks in Economics	25	28	35	32	31	36	29	38	34	32
Marks in Statistics	43	46	49	41	36	32	31	30	33	39

Q6: Under an employment promotion programme, it is proposed to allow sale of newspapers on the buses during off-peak hours. The vendor can purchase the newspapers at a special concessional rate of 25 paise per copy against the selling price of 40 paise. Any unsold copies are however, a dead loss. A vendor has estimated the following probability distribution for the number of copies demanded:

Number of copies	15	16	17	18	19	20
Probability	0.04	0.19	0.33	0.26	0.11	0.07

How many copies should he order so that his expected profits will be maximum?

IV. BINOMIAL COEFFICIENTS

n	$\binom{n}{0}$	$\binom{n}{1}$	$\binom{n}{2}$	$\binom{n}{3}$	$\binom{n}{4}$	$\binom{n}{5}$	$\binom{n}{6}$	$\binom{n}{7}$	$\binom{n}{8}$	$\binom{n}{9}$	$\binom{n}{10}$
0	1										
1	1	1									
2	1	2	1								
3	1	3	3	1							
4	1	4	6	4	1						
5	1	5	10	10	5	1					
6	1	6	15	20	15	6	1				
7	1	7	21	35	35	21	7	1			
8	1	8	28	56	70	56	28	8	1		
9	1	9	36	84	126	126	84	36	9	1	
10	1	10	45	120	210	252	210	120	45	10	1
11	1	11	55	165	330	462	462	330	165	55	11
12	1	12	66	220	495	792	924	792	495	220	66
13	1	13	78	286	715	1287	1716	1716	1287	715	286
14	1	14	91	364	1001	2002	3003	3003	2002	1001	364
15	1	15	105	455	1365	3003	5005	6435	6435	3005	455
16	1	16	120	560	1820	4368	8008	11440	12870	11440	8008
17	1	17	136	680	2380	6188	12376	19448	24310	24310	19448
18	1	18	153	816	3060	8568	18564	31824	43758	48620	43758
19	1	19	171	969	3876	11628	27132	50388	75582	92378	92378
20	1	20	190	1140	4845	15504	38760	77520	125970	167960	184756

V. VALUES OF e^{-m} (For Computing Poisson Probabilities)
($0 < m < 1$)

m	0	1	2	3	4	5	6	7	8	9
0.0	1.0000	.9900	.9802	.9704	.9608	.9512	.9418	.9324	.9231	.9139
0.1	0.9048	.8958	.8860	.8781	.8694	.8607	.8521	.8437	.8353	.8270
0.2	0.8187	.8108	.8025	.7945	.7866	.7788	.7711	.7634	.7558	.7483
0.3	0.7408	.7334	.7261	.7189	.7118	.7047	.6977	.6907	.6839	.6771
0.4	0.6703	.6636	.6570	.6505	.6440	.6376	.6313	.6250	.6188	.6126
0.5	0.6065	.6005	.5945	.5886	.5827	.5770	.5712	.5655	.5599	.5543
0.6	0.5488	.5434	.5379	.5326	.5278	.5220	.5160	.5117	.5066	.5016
0.7	0.4966	.4916	.4868	.4810	.4771	.4724	.4670	.4630	.4584	.4538
0.8	0.4493	.4449	.4404	.4360	.4317	.4274	.4232	.4190	.4148	.4107
0.9	0.4066	.4025	.3985	.3946	.3906	.3867	.3829	.3791	.3753	.3716

(m = 1, 2, 3, ..., 10)

m	1	2	3	4	5	6	7	8	9	10
e^{-m}	.36788	.13534	.04979	.01832	.00698	.00278	.00092	.00035	.00012	.00004

Note: To obtain values of e^{-m} for other values of m , use the laws of exponents.

Example. $e^{-2.35} = (e^{-2.00})(e^{-0.35}) = (.13534)(.7047) = .095374$

X. 5% POINTS OF FISHER'S F-DISTRIBUTION

APPENDIX

m \ n	1	2	3	4	5	6	7	8	9	10	12	15	20	30	60	∞
1	161.45	199.50	215.71	224.58	230.16	233.99	236.77	238.88	240.54	241.88	243.91	245.95	248.01	250.09	252.20	254.32
2	18.513	19.000	19.164	19.247	19.296	19.330	19.353	19.371	19.385	19.396	19.413	19.420	19.446	19.462	19.479	19.496
3	10.128	9.5521	9.2706	9.1172	9.0135	8.9408	8.8908	8.8452	8.8123	8.7855	8.7446	8.7029	8.6602	8.6166	8.5720	8.5265
4	7.7092	6.9443	6.5914	6.3883	6.2560	6.1631	6.0942	6.0410	5.9988	5.9644	5.9117	5.8578	5.8025	5.7459	5.6878	5.6281
5	6.6079	5.7861	5.4095	5.1922	5.0503	4.9503	4.8753	4.8183	4.7725	4.7351	4.6777	4.6188	4.5581	4.4957	4.4314	4.3650
6	5.9874	5.1433	4.7571	4.5337	4.3874	4.2839	4.2066	4.1468	4.0990	4.0600	3.9999	3.9381	3.8742	3.8082	3.7398	3.6688
7	5.5914	4.7374	4.3468	4.1203	3.9715	3.8660	3.7870	3.7257	3.6767	3.6365	3.5747	3.5108	3.4445	3.3758	3.3043	3.2298
8	5.3177	4.4590	4.0662	3.8378	3.6875	3.5806	3.5005	3.4381	3.3881	3.3472	3.2840	3.2184	3.1503	3.0794	3.0050	2.9278
9	5.1174	4.2565	3.8626	3.6331	3.4817	3.3738	3.2927	3.2296	3.1789	3.1373	3.0729	3.0001	2.9365	2.8637	2.7922	2.7067
10	4.9646	4.1028	3.7083	3.4780	3.3258	3.2172	3.1355	3.0717	3.0204	2.9782	2.9130	2.8450	2.7740	2.6996	2.6211	2.5379
11	4.8443	3.9823	3.5874	3.3567	3.2039	3.0946	3.0123	2.9480	2.8962	2.8536	2.7876	2.7186	2.6464	2.5705	2.4901	2.4045
12	4.7272	3.8653	3.4703	3.2392	3.0861	2.9761	2.8934	2.8291	2.7764	2.7334	2.6666	2.5966	2.5233	2.4463	2.3642	2.2762
13	4.6672	3.8056	3.4105	3.1791	3.0254	2.9153	2.8321	2.7669	2.7144	2.6710	2.6037	2.5331	2.4589	2.3803	2.2966	2.2064
14	4.6001	3.7389	3.3439	3.1122	2.9582	2.8477	2.7642	2.6987	2.6458	2.6021	2.5342	2.4630	2.3879	2.3082	2.2230	2.1307
15	4.5431	3.6823	3.2874	3.0556	2.9013	2.7905	2.7066	2.6408	2.5876	2.5437	2.4753	2.4035	2.3275	2.2468	2.1601	2.0658
16	4.4940	3.6337	3.2389	3.0069	2.8524	2.7413	2.6572	2.5911	2.5377	2.4935	2.4247	2.3522	2.2756	2.1938	2.1058	2.0095
17	4.4513	3.5915	3.1968	2.9647	2.8100	2.6987	2.6143	2.5480	2.4943	2.4499	2.3807	2.3077	2.2304	2.1477	2.0581	1.9604
18	4.4139	3.5546	3.1599	2.9277	2.7729	2.6613	2.5767	2.5102	2.4563	2.4117	2.3421	2.2686	2.1906	2.1071	2.0166	1.9168
19	4.3808	3.5219	3.1274	2.8951	2.7401	2.6283	2.5435	2.4768	2.4227	2.3779	2.3080	2.2341	2.1555	2.0712	1.9796	1.8790
20	4.3513	3.4928	3.0984	2.8661	2.7100	2.5980	2.5140	2.4471	2.3928	2.3479	2.2776	2.2033	2.1242	2.0391	1.9464	1.8432
21	4.3248	3.4668	3.0725	2.8401	2.6840	2.5727	2.4876	2.4205	2.3661	2.3210	2.2504	2.1767	2.0960	2.0102	1.9165	1.8117
22	4.3009	3.4434	3.0491	2.8167	2.6603	2.5487	2.4638	2.3965	2.3419	2.2967	2.2258	2.1508	2.0707	1.9842	1.8895	1.7831
23	4.2793	3.4221	3.0280	2.7955	2.6390	2.5277	2.4422	2.3748	2.3201	2.2747	2.2036	2.1282	2.0476	1.9605	1.8649	1.7570
24	4.2597	3.4026	3.0085	2.7763	2.6207	2.5092	2.4238	2.3561	2.3002	2.2547	2.1834	2.1077	2.0267	1.9390	1.8424	1.7331
25	4.2417	3.3852	2.9912	2.7587	2.6030	2.4914	2.4057	2.3377	2.2811	2.2355	2.1649	2.0889	2.0075	1.9192	1.8217	1.7110
26	4.2252	3.3690	2.9751	2.7426	2.5868	2.4751	2.3893	2.3205	2.2635	2.2177	2.1479	2.0716	1.9898	1.9010	1.8027	1.6906
27	4.2100	3.3541	2.9604	2.7278	2.5719	2.4591	2.3732	2.3053	2.2481	2.2023	2.1323	2.0558	1.9736	1.8842	1.7851	1.6717
28	4.1960	3.3404	2.9467	2.7141	2.5581	2.4453	2.3593	2.2913	2.2340	2.1880	2.1179	2.0411	1.9586	1.8687	1.7689	1.6541
29	4.1830	3.3277	2.9340	2.7014	2.5454	2.4324	2.3463	2.2782	2.2209	2.1748	2.1045	2.0275	1.9446	1.8543	1.7537	1.6377
30	4.1709	3.3158	2.9223	2.6896	2.5336	2.4205	2.3343	2.2662	2.2087	2.1626	2.0921	2.0148	1.9317	1.8409	1.7396	1.6223
40	4.0648	3.2317	2.8387	2.6060	2.4495	2.3359	2.2490	2.1802	2.1240	2.0772	2.0053	1.9245	1.8399	1.7444	1.6373	1.5089
60	4.0012	3.1504	2.7581	2.5252	2.3688	2.2540	2.1665	2.0970	2.0401	1.9928	1.9174	1.8364	1.7480	1.6491	1.5343	1.3803
120	3.9201	3.0718	2.6802	2.4472	2.2900	2.1750	2.0867	2.0164	1.9588	1.9105	1.8337	1.7505	1.6587	1.5543	1.4290	1.2509
∞	3.8415	2.9957	2.6049	2.3719	2.2141	2.0988	2.0096	1.9384	1.8799	1.8307	1.7522	1.6664	1.5705	1.4591	1.3180	1.0000

For $m > 10$ interpolate using 60/ m
 For $n > 30$ interpolate using 120/ n

ANNEXURE 3

Degrees of Freedom	Level of Significance		
	10%	5%	1%
1	2.706	3.841	6.635
2	4.605	5.991	9.210
3	6.251	7.815	11.345
4	7.779	9.488	13.277
5	9.236	11.071	15.086
6	10.645	12.592	16.812
7	12.017	14.067	18.475
8	13.362	15.507	20.090
9	14.684	16.919	21.666
10	15.987	18.307	23.209
11	17.275	19.675	24.725
12	18.549	21.026	26.217
13	19.812	22.362	27.688
14	21.064	23.685	29.141
15	22.307	24.996	30.578
16	23.542	26.296	32.000
17	24.769	27.587	33.409
18	25.989	28.869	34.805
19	27.204	30.144	36.191
20	28.412	31.410	37.566
21	29.615	32.671	38.932
22	30.813	33.924	40.289
23	33.007	35.172	41.638
24	33.196	36.415	42.980
26	34.382	37.652	44.314
26	35.363	38.885	45.642
27	36.741	40.113	46.963
28	37.916	41.337	48.278
29	38.087	42.587	49.588
30	40.256	43.773	50.892

Note: For degrees of freedom greater than 30, the quantity $\sqrt{2x^2 - \sqrt{2v - 1}}$ may be used as a normal variate with unit variance

Annexures 1-4

ANNEXURE 1

Z	0.00	0.01	0.02	0.03	0.04	0.05	0.06	0.07	0.08	0.09
0.0	0.0000	0.0040	0.0080	0.0120	0.0160	0.0199	0.0239	0.0279	0.0319	0.0359
0.1	0.0398	0.0438	0.0478	0.0517	0.0557	0.0596	0.0636	0.0675	0.0714	0.0753
0.2	0.0793	0.0832	0.0871	0.0910	0.0948	0.0987	0.1026	0.1064	0.1103	0.1141
0.3	0.1179	0.1217	0.1255	0.1293	0.1331	0.1368	0.1406	0.1443	0.1480	0.1517
0.4	0.1554	0.1591	0.1628	0.1664	0.1700	0.1736	0.1772	0.1808	0.1844	0.1879
0.5	0.1915	0.1950	0.1985	0.2019	0.2054	0.2088	0.2123	0.2157	0.2190	0.2224
0.6	0.2257	0.2291	0.2324	0.2357	0.2389	0.2422	0.2454	0.2486	0.2517	0.2548
0.7	0.2580	0.2611	0.2642	0.2673	0.2704	0.2734	0.2764	0.2794	0.2823	0.2852
0.8	0.2881	0.2910	0.2939	0.2967	0.2995	0.3023	0.3051	0.3078	0.3106	0.3133
0.9	0.3159	0.3186	0.3212	0.3238	0.3264	0.3289	0.3315	0.3340	0.3365	0.3389
1.0	0.3413	0.3438	0.3461	0.3485	0.3508	0.3531	0.3554	0.3577	0.3599	0.3621
1.1	0.3643	0.3665	0.3686	0.3708	0.3729	0.3749	0.3770	0.3790	0.3810	0.3830
1.2	0.3849	0.3869	0.3888	0.3907	0.3925	0.3944	0.3962	0.3980	0.3997	0.4015
1.3	0.4032	0.4049	0.4066	0.4082	0.4099	0.4115	0.4131	0.4147	0.4162	0.4177
1.4	0.4192	0.4207	0.4222	0.4236	0.4251	0.4265	0.4279	0.4292	0.4306	0.4319
1.5	0.4332	0.4345	0.4357	0.4370	0.4382	0.4394	0.4406	0.4418	0.4429	0.4441
1.6	0.4452	0.4463	0.4474	0.4484	0.4495	0.4505	0.4515	0.4525	0.4535	0.4545
1.7	0.4554	0.4564	0.4573	0.4582	0.4591	0.4599	0.4608	0.4616	0.4625	0.4633
1.8	0.4641	0.4649	0.4656	0.4664	0.4671	0.4678	0.4686	0.4693	0.4699	0.4706
1.9	0.4713	0.4719	0.4726	0.4732	0.4738	0.4744	0.4750	0.4756	0.4761	0.4767
2.0	0.4772	0.4778	0.4783	0.4788	0.4793	0.4798	0.4803	0.4808	0.4812	0.4817
2.1	0.4821	0.4826	0.4830	0.4834	0.4838	0.4842	0.4846	0.4850	0.4854	0.4857
2.2	0.4861	0.4864	0.4868	0.4871	0.4875	0.4878	0.4881	0.4884	0.4887	0.4890
2.3	0.4893	0.4896	0.4898	0.4901	0.4804	0.4906	0.4909	0.4911	0.4913	0.4916
2.4	0.4918	0.4920	0.4922	0.4925	0.4927	0.4929	0.4931	0.4932	0.4934	0.4936
2.6	0.4953	0.4955	0.4956	0.4957	0.4959	0.4960	0.4961	0.4962	0.4963	0.4964
2.8	0.4974	0.4975	0.4976	0.4977	0.4977	0.4978	0.4979	0.4979	0.4980	0.4981
2.9	0.4981	0.4982	0.4982	0.4983	0.4984	0.4984	0.4985	0.4985	0.4986	0.4986
3.0	0.4987	0.4987	0.4987	0.4988	0.4988	0.4989	0.4989	0.4989	0.4990	0.4990

ANNEXURE 2

Degrees of Freedom	Level of Significance		
	1%	5%	10%
1	63.657	12.706	6.314
2	9.925	4.303	2.920
3	5.841	3.182	2.353
4	4.604	2.776	2.132
5	4.032	2.571	2.015
6	3.707	2.447	1.943
7	3.499	2.365	1.895
8	3.355	2.306	1.860
9	3.250	2.262	1.833
10	3.169	2.228	1.812
11	3.106	2.201	1.796
12	3.055	2.179	1.782
13	3.012	2.160	1.771
14	2.977	2.145	1.761
15	2.947	2.131	1.753
16	2.921	2.120	1.746
17	2.898	2.110	1.740
18	2.878	2.101	1.734
19	2.861	2.093	1.729
20	2.846	2.086	1.725
21	2.831	2.080	1.721
22	2.819	2.074	1.717
23	2.807	2.069	1.714
24	2.797	2.064	1.711
25	2.787	2.060	1.708
26	2.779	2.056	1.706
27	2.771	2.052	1.703
28	2.763	2.048	1.701
29	2.756	2.045	1.699
α	2.576	1.960	1.645

Note: These table values of t are in respect of two-tailed tests. If we use the t -distribution for one-tailed test then we are interested in determining the area located in one tail. So to find the appropriate t -value for a one-tailed test say at a 5% level with 12 degrees of freedom, then we should look in the above table under the 10% column opposite the 12 degrees of freedom row. (This value will be 1.782). This is true because the 10% column represents 10% of the area under the curve contained in both tails combined, and so it also represents 5% of the area under the curve contained in each of the tails separately.